

Parasites of the Potato Tuberworm, Tomato Pinworm, and Other, Closely Related Gelechiids

EARL R. OATMAN¹ and GARY R. PLATNER¹

ABSTRACT

Parasites reared from the potato tuberworm, tomato pinworm, and other closely related gelechiids in the Americas are presented. The parasite complex associated with each gelechiid and the area of origin of the potato tuberworm and tomato pinworm are discussed briefly.

The potato tuberworm, *Phthorimaea operculella* (Zeller), is a worldwide pest of potato, tomato, eggplant, tobacco, and other Solanaceae. It is especially serious in dry, subtropical climates and during dry seasons in areas normally having adequate rainfall (Lloyd, 1972). The potato tuberworm occurs in all major potato-growing areas of central and southern California, but not in the extreme northern part of the state (Bacon, 1960). In southern California, where two irrigated crops of potatoes are grown annually, the pest is active throughout the year (Oatman & Platner, 1974).

The cultivated potato, *Solanum tuberosum* L., is the most common host of the potato tuberworm. The pest larvae mine the foliage, stems, and tubers. Most of the economic damage is to the exposed tubers prior to harvest; up to 100% may be infested if preventive measures, such as deep planting and ridging, are not used (Shelton & Wyman, 1979).

In the United States, the tomato pinworm, *Keiferia lycopersicella* (Walsingham), is a serious economic pest of fresh market tomatoes in California (Oatman, 1970), Florida (Wolfenbarger & Poe, 1973), and Texas (Harding, 1971). Based on personal observations by the senior author, it also is a serious pest of tomatoes in Hawaii and commonly infests commercial eggplant in southern California.

Because of the economic importance of potatoes and tomatoes in California's agricultural economy and the severity of the potato tuberworm and tomato pinworm to these crops, studies were initiated in 1963 in southern California to determine the parasite complexes associated with both pests on their respective host plants (Oatman, 1970; Oatman & Platner, 1974; Oatman et al., 1979). During the course of these studies, foreign exploration trips were made to potato- and tomato-growing regions of Mexico, Central, and South America by the senior author to attempt to delineate the area of origin of the potato tuberworm and tomato pinworm and to collect

¹Division of Biological Control, Department of Entomology, University of California, Riverside, California 92521 U.S.A.

exotic parasites of each for release in southern California. Two such trips were made to Mexico (February, 1967 and 1981), two to Costa Rica (February, 1967, and April, 1973), and one each to Colombia (May, 1973) and Guatemala (February, 1977). Additionally, trips were made to Hawaii (June-December, 1977) and the Lower Rio Grande Valley, Texas (June, 1979) to collect parasites of the tomato pinworm. In all cases, parasitized material was collected from unsprayed plants, usually growing individually or in small plots, and shipped by air to the Division of Biological Control, University of California, Riverside, where the material was processed through the Division's quarantine facility. Parasites thus collected are shown in Tables 1 and 2. Included in Table 2 are parasites of the tomato pinworm collected in 1977 by Mr. Robert Burkhardt, Division of Plant Industry, Department of Agriculture, State of Hawaii, during a foreign exploration trip to Arizona, California, Texas, Mexico, Guatemala, Honduras, and El Salvador.

From 1963 through 1965, the Department of Biological Control, University of California, Riverside, cooperated with the Commonwealth Institute of Biological Control (CIBC) by handling shipments of parasites of the potato tuberworm through the Department's quarantine facility. CIBC field agents collected parasitized material from potato foliage during the summers of 1962-63 and 1963-64 in Argentina and Uruguay and during the summer of 1964-65 in Peru (Lloyd, 1972). Parasites received from these shipments, plus five additional ones from CIBC facilities in South Africa and India, are listed in Table 1. Except for *Microplitis minutalis* Muesebeck, *Rhetisymphiesis phthorimaeae* Blanchard and two species of *Apanteles* from Peru, sufficient numbers of each parasite species were obtained to establish insectary cultures. All the cultured species were released against the potato tuberworm in southern California (Oatman & Platner, 1974). Concurrently, cultures of the South American parasites were shipped to the Indian Station, CIBC, Bangalore, India, for propagation and distribution to other Commonwealth countries participating in the CIBC project on biological control of the potato tuberworm (Lloyd, 1972). Additional species (eight insects and one nematode), reported as parasites of the potato tuberworm in South America (Table 1), were not collected.

Of the ten parasites received from CIBC and released in southern California, only *Copidosoma koehleri* (Blanchard) and *Orgilus lepidus* Muesebeck were established (Oatman & Platner, 1974). Results of general surveys for parasites of the potato tuberworm in 1963 and 1964 and selective surveys in 1968 and 1969 showed that 17 primary parasites are present in southern California (Table 1), at least 15 being indigenous. *Copidosoma koehleri* was not recovered in the 1968-69 surveys and, thus, may have failed to survive.

Parasites of the potato tuberworm, collected in Mexico (ca. 100 miles north of Mexico City), were the same species as those in southern California. Of the seven parasites collected in Costa Rica (near Cartago), only one ichneumonid and two eulophids were in genera different from those collected in California (Table 1).

TABLE 1. Parasites reared from *Phthorimaea operculella* on potato.

Parasite	Collected in
Braconidae:	
<i>Agathis gibbosa</i> (Say)	California
* <i>Agathis tandilensis</i> Blanchard	Argentina
* <i>Agathis unicolor</i> (Schrottky)	Argentina
<i>Apanteles dignus</i> Muesebeck	California
<i>Apanteles scutellaris</i> Muesebeck	California, Mexico
* <i>Apanteles subandinus</i> Blanchard	Argentina, Peru
* <i>Apanteles</i> spp. (2) ^a	Peru
* <i>Bracon cayanus</i> Blanchard	Argentina
<i>Bracon gelechiae</i> Ashmead	California
* <i>Chelonus curvumaculatus</i> Cameron	S. Africa
<i>Chelonus kellyae</i> Marsh	Costa Rica
<i>Chelonus phthorimaeae</i> Gahan	California, Mexico
* <i>Microplitis minuralis</i> Muesebeck	Uruguay
<i>Ongilus jenniae</i> Marsh	Costa Rica
<i>Ongilus lateralis</i> (Cresson)	California
* <i>Ongilus lepidus</i> Muesebeck	California, Argentina
* <i>Ongilus parvus</i> Turner	S. Africa
<i>Ongilus</i> spp.	California, Costa Rica
* <i>Rogus</i> spp.	S. America
Ichneumonidae:	
* <i>Campoplex haywardi</i> Blanchard	Argentina, Uruguay
<i>Campoplex phthorimaeae</i> (Cushman)	California, Mexico
<i>Diadegma blackburni</i> (Cameron)	California, Hawaii
<i>Diadegma compressus</i> (Cresson)	California
* <i>Diadegma stellenboschense</i> (Cameron)	S. Africa
<i>Diadegma</i> sp.	California
<i>Nephera fuscifemora</i> (Graf)	California
* <i>Nythobia</i> spp. (2) ^a	Cyprus, India
<i>Pristomerus spinator</i> (F.)	California
* <i>Temelucha</i> spp.	California, Argentina, Costa Rica
<i>Trathala</i> sp.	Costa Rica
Encyrtidae:	
* <i>Apsilophrys oeceticola</i> (DeSantis)	Peru
* <i>Copidosoma koehleri</i> (Blanchard)	Argentina, Brazil, Chile, Peru
Eulophidae:	
<i>Elachertus</i> sp.	Costa Rica
<i>Hyssopus</i> sp.	Costa Rica
* <i>Rhetisymphiesis phthorimaeae</i> Blanchard	Argentina, Peru
<i>Symphiesis stigmatipennis</i> Girault	California

^aTwo species.

*Received in quarantine (1963-1967), Department of Biological Control, University of California, Riverside, from Commonwealth Institute of Biological Control (CIBC).

**Reported as parasites of *Phthorimaea operculella* in South America by CIBC.

TABLE 1. Parasites reared from *Phthorimaea operculella* on potato. (Continued)

Parasite	Collected in
Pteromalidae:	
** <i>Lophocomodia</i> sp.	Argentina, Peru
Scelionidae:	
** <i>Telenomus</i> sp.	S. America
Trichogrammatidae:	
<i>Trichogramma pretiosum</i> Riley	California
Tachinidae:	
** <i>Incarnia cuzcensis</i> Townsend	Peru
Mermithidae:	
** <i>Hexameris</i> sp.	Argentina

**Reported as parasites of *Phthorimaea operculella* in South America by CIBC.

TABLE 2. Parasites reared from *Keiferia lycopersicella* on tomato.

Parasite	Collected in ^a
Braconidae:	
<i>Agathis</i> sp.	TX
<i>Apanteles dignus</i> Muesebeck	AZ, CA, HI, TX, EL SALV., GUAT., HOND., MEX.
<i>Apanteles scutellaris</i> Muesebeck	CA, HI, TX, MEX.
<i>Bracon gelechiae</i> Ashmead	CA
<i>Bracon</i> spp.	TX, GUAT., MEX.
<i>Chelonus blackburni</i> Cameron	HI
<i>Chelonus phthorimaeae</i> Gahan	AZ, CA, TX, GUAT., MEX.
<i>Orgilus</i> spp.	CA, TX, GUAT., MEX.
<i>Pavahornius pallidipes</i> (Ashmead)	AZ, CA, TX, GUAT., MEX.
Ichneumonidae:	
<i>Campoplex phthorimaeae</i> (Cushman)	CA
* <i>Campoplex</i> n. sp.	TX, MEX.
<i>Pristomerus hawaiiensis</i> Perkins	HI
<i>Pristomerus spinator</i> (F.)	HI, TX, GUAT., MEX.
<i>Trathala flavoorbitalis</i> Cameron	HI
Eulophidae:	
* <i>Elasmus nigripes</i> Howard	TX, GUAT., HOND., MEX.
<i>Sympiesis stigmatipennis</i> Girault	CA
Pteromalidae:	
* <i>Zatropis</i> sp. (near <i>Tortricidis</i>)	GUAT.
Bethylidae:	
<i>Goniozus</i> sp. (near <i>Platynotae</i>)	CA, GUAT.
Trichogrammatidae:	
<i>Trichogramma pretiosum</i> Riley	CA

^aAZ = Arizona; CA = California; HI = Hawaii; TX = Texas; EL SALV. = El Salvador; GUAT. = Guatemala; HOND. = Honduras; MEX. = Mexico.

*Species collected by Burkhart that differed from those collected by Oatman.

The parasite complexes in North and South America are similar (Table 1), suggesting the existence of ecological homologues on the two continents. As noted by Lloyd (1972), parasites occurring most commonly in each continent include one egg-larval parasite each, two gregarious external larval parasites each, six solitary, internal larval parasites (Braconidae) each, and three solitary, internal larval parasites (Ichneumonidae) each. The biologies of most of these have been published (Leon & Oatman, 1968; Oatman et al., 1969; Djamin, 1970; Cardona & Oatman, 1971; Odebiyi & Oatman, 1972; Oatman & Platner, 1974; Cardona & Oatman, 1975; Odebiyi & Oatman, 1977; Flanders & Oatman, 1982; Teran, 1983; Powers & Oatman, 1985).

Twenty-one species of primary parasites of the tomato pinworm have been recorded (Table 2). Ten of these have been recorded from southern California, nine each from Guatemala, Mexico and Texas, and six from Hawaii. Of the ten parasites recorded from southern California, six are common (Oatman, 1970; Oatman et al., 1979). Overall, *Apanteles dignus* Muesebeck is the predominant parasite, having been found in all areas surveyed. Except for those from Hawaii, all the parasites (Table 2) occur naturally in the area(s) from which they were collected. Based on the results of foreign exploration trips and personal observations, the tomato pinworm does not occur in Costa Rica, Argentina, or Colombia. The available evidence strongly indicates that the tomato pinworm originated in the area that is now Mexico and Guatemala. The large numbers of parasites (Table 2) associated with this pest species support this. Also, the tomato (principal host) is thought to have originated somewhere in Central America. The higher number of parasites in southern California probably is due to the more extensive studies conducted there (Oatman, 1970; Oatman et al., 1979). The greatest number of different primary parasites (seven) collected by Burkhart in 1977 was near Sanarate, Guatemala.

Neither the potato tuberworm nor the tomato pinworm was collected during the senior author's foreign exploration trip to Colombia in 1973. All the gelechiids collected from both potato and tomato in the commercial-growing areas were *Scrobipalpus absoluta* (Meyrick). In Colombia, this gelechiid mines the leaves of potato, but does not infest the tubers. In tomato, *S. absoluta* mines the leaves and infests the fruit. It is a serious economic pest in that it infests both green and ripe tomatoes, injuring up to 100% of the fruit in spite of repeated applications of insecticides (personal observations). Fifteen parasites were reared from the *S. absoluta*-infested material collected in Colombia in 1973 (Table 3).

While collecting *S. absoluta* near Palmira, Colombia, gelechiid-mined leaves from a native solanaceous weed, *Solanum saponaceum* Dun., also were collected and shipped to Riverside. The gelechiid involved was identified as *Scrobipalpus isoclorus* Povolny. In 1975, Mr. Alfredo Saldarriaga, Programa Entomologia, Instituto Colombiano Agropecuario, Palmira, Colombia, shipped additional parasites of *S. isoclorus* to us for identification. These species also are listed in Table 3. As both *S. absoluta* and *S. isoclorus* are closely related to the potato tuberworm and the tomato pinworm, their parasite

TABLE 3: Parasites reared from *Scrobipalpula absoluta* on potato and tomato in Colombia and from *Scrobipalpula isoclera* on *Solanum saponaceum* in the Cauca Valley, Palmira, Colombia.

<i>S. absoluta</i>	<i>S. isoclera</i>
Bethylidae:	
<i>Goniozus nigrifemur</i> (Ashmead)	<i>Goniozus</i> sp.
Braconidae:	
<i>Apanteles dignus</i> Muesebeck	<i>Apanteles epinotae</i> Vicrick
<i>Apanteles gelechiidivorus</i> Marsh	<i>Apanteles oatmani</i> Marsh
<i>Apanteles</i> sp.	<i>Apanteles</i> sp.
<i>Bracon</i> sp.	<i>Bracon lucileae</i> Marsh
	<i>Bracon</i> sp.
	<i>Chelonus (Microchelonus) johni</i> Marsh
	<i>Mirax malcomi</i> Marsh
Chalcididae:	
<i>Invreia</i> sp.	<i>Brachymeria</i> sp. ^a
	<i>Spilochalcis hirtifemora</i> (Ashmead) ^a
Eulophidae:	
<i>Elasmus</i> sp.	<i>Proacrias</i> sp.
<i>Sympiesis</i> sp.	<i>Tetrastichus</i> nr. <i>chapadae</i> Ashmead
<i>Tetrastichus</i> sp.	<i>Tetrastichus</i> sp.
	<i>Zagrammosoma</i> spp.
Eupelmidae:	
<i>Anastatus</i> sp.	
Ichneumonidae:	
<i>Diadegma</i> sp.	<i>Pristomerus</i> spp.
<i>Pristomerus</i> sp.	
<i>Temelucha</i> sp.	
Perilampidae:	
	<i>Perilampus</i> sp. ^a
Pteromalidae:	
	<i>Zatropus</i> sp.
Trichogrammatidae:	
<i>Trichogramma pretiosum</i> Riley	
Tachinidae:	
<i>Elfia</i> (= <i>Plectops</i>) sp.	

^aProbably hyperparasites.

complexes should be considered in planning an introduction program for biological control of the last two pests. Altogether, about fifteen primary parasites were reared from each of *S. absoluta* and *S. isoclorata* (Table 3). The *Apanteles* sp., *Bracon* sp., *Goniozus* sp., *Tetrastichus* sp., and *Pristomerus* sp. listed probably are the same species on both gelechiids. Nevertheless, about 25 different primary parasites were recorded altogether.

Lloyd (1972) stated that "there is good evidence for the claim that the cultivated potato originated in South America and it is a reasonable presumption that the tuber moth also spread over the world from that same general area." Such a statement helps little in delineating the pest's area of origin, considering the size of the land area (continent). Based on personal observations, the potato tuberworm does not occur in Colombia (at least not in 1973), but is present in Venezuela, where it is a pest on tomato near Barquisimeto. Its absence from Colombia is difficult to understand if, in fact, it originated in South America.

Thus, it is equally reasonable to presume that the potato tuberworm originated in North America, as both the pest and its principal hosts, both wild (e.g., *Datura* spp.) and cultivated, commonly occur here (personal observations). Lloyd (1972) noted that several early workers, concerned with the potato tuberworm, felt that the pest originated in North America. The large parasite complex associated with the potato tuberworm in southern California, and the occurrence of the same parasite species in south-central Mexico, suggest that the pest has been here equally as long as it has been in South America. Certainly, those interested in collecting parasites of the potato tuberworm would do well to concentrate their search in the drier, subtropical areas of North America.

ACKNOWLEDGMENTS

Thanks are extended to B. D. Burks, R. W. Carlson, P. M. Marsh, A. S. Menke, C. F. W. Muesebeck, C. W. Sabrosky and L. M. Walkley, Systematic Entomology Laboratory, Insect Identification and Beneficial Insect Introduction Institute (IIBIII), USDA, Beltsville, Maryland, and G. Gordh and J. D. Pinto, Division of Biological Control, University of California, Riverside, for identifying the parasites, and to R. W. Hodges and D. M. Weisman, Systematic Entomology Laboratory, IIBIII, for identifying the Gelechiidae. We also thank Po-Yung Lai, Division of Plant Industry, Department of Agriculture, State of Hawaii, for permission to include parasites of the tomato pinworm, collected by R. Burkhart, and to A. Saldarriaga, Programa Entomologia, Instituto Colombiano Agropecuario, for collecting additional parasites of *Scrobipalpus isoclorata*.

REFERENCES CITED

- Bacon, O. G. 1960. Control of the potato tuberworm in potatoes. J. Econ. Entomol. 53:853-871.
- Cardona, C. and E. R. Oatman. 1971. Biology of *Apanteles dignus* (Hymenoptera: Braconidae), a primary parasite of the tomato pinworm. Ann. Entomol. Soc. Amer. 64:996-1007.
- Cardona, C. and E. R. Oatman. 1975. Biology and physical ecology of *Apanteles subandinus* Blanchard (Hymenoptera: Braconidae), with notes on temperature responses of *Apanteles scutellaris* Muesebeck and its host, the potato tuberworm. Hilgardia 43:1-51.
- Djain, A. 1970. Biologies of the tomato pinworm, *Keiferia lycopersicella* (Walsingham) (Lepidoptera: Gelechiidae) and its parasite, *Apanteles scutellaris* Muesebeck (Hymenoptera: Braconidae), with special reference to the influence of temperature on population increase. Ph.D. Dissertation, Univ. Calif., Berkeley. Unpublished. 119 pp.
- Flanders, R. V. and E. R. Oatman. 1982. Laboratory studies on the biology of *Orgilus jenniae* (Hymenoptera: Braconidae), a parasitoid of the potato tuberworm, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). Hilgardia 50:1-33.
- Harding, J. A. 1971. Field comparisons of insecticidal sprays for control of four tomato insects in south Texas. J. Econ. Entomol. 64:1302-1304.
- Leon, J. K. L. and E. R. Oatman. 1968. Biology of *Campoplex haywardi* (Hymenoptera: Ichneumonidae), a primary parasite of the potato tuberworm. Ann. Entomol. Soc. Amer. 61:26-36.
- Lloyd, D. C. 1972. Some South American parasites of the potato tuber moth, *Phthorimaea operculella* (Zeller) and remarks on those in other continents. Commonwealth Inst. Bio. Control, Commonwealth Agric. Bureau, Tech. Bull. 15:35-49.
- Oatman, E. R. 1970. Ecological studies of the tomato pinworm on tomato in southern California. J. Econ. Entomol. 63:1531-1534.
- Oatman, E. R. and G. R. Platner. 1974. Biology of *Temelucha* sp., *Platensis* group (Hymenoptera: Ichneumonidae), a primary parasite of the potato tuberworm. Ann. Entomol. Soc. Amer. 67:275-280.
- Oatman, E. R. and G. R. Platner. 1974. Parasitization of the potato tuberworm in southern California. Environ. Entomol. 3:262-264.
- Oatman, E. R., G. R. Platner, and P. D. Greany. 1969. Biology of *Orgilus lepidus* (Hymenoptera: Braconidae), a primary parasite of the potato tuberworm. Ann. Entomol. Soc. Amer. 62:1407-1414.
- Oatman, E. R., J. A. Wyman, and G. R. Platner. 1979. Seasonal occurrence and parasitization of the tomato pinworm on fresh market tomatoes in southern California. Environ. Entomol. 8:661-664.
- Odebiyi, J. A. and E. R. Oatman. 1972. Biology of *Agathis gibbosa* (Hymenoptera: Braconidae), a primary parasite of the potato tuberworm. Ann. Entomol. Soc. Amer. 64:1104-1114.
- Odebiyi, J. A. and E. R. Oatman. 1977. Biology of *Agathis unicolor* (Schrottky) and *Agathis gibbosa* (Say) (Hymenoptera: Braconidae), primary parasites of the potato tuberworm. Hilgardia 45:123-151.
- Powers, N. R. and E. R. Oatman. 1985. Biology and temperature responses of *Chelonus kelliae* and *Chelonus phthorimaeae* (Hymenoptera: Braconidae) and their host, the potato tuberworm, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). Hilgardia 52:1-32.
- Shelton, A. M. and J. A. Wyman. 1979. Potato tuberworm damage to potatoes under different irrigation and cultural practices. J. Econ. Entomol. 72:261-264.
- Teran, J. B. 1983. Comparative biologies of *Parahormius pallididipes* (Ashmead) (Hymenoptera: Braconidae) and *Sympiesis stigmatipennis* Girault (Hymenoptera: Eulophidae), ectoparasites of the tomato pinworm, *Keiferia lycopersicella* (Walsingham) (Lepidoptera: Gelechiidae). Ph.D. Dissertation, Univ. Calif., Riverside. Unpublished. 147 pp.
- Wolfenbarger, D. O. and S. L. Poe. 1973. Tomato pinworm control. Proc. Fla. St. Hortic. Soc. 86:139-143.